

* doi:10.4317/medoral.17644048

<http://dx.doi.org/10.4317/medoral.17644048>

Introduction: Oral bisphosphonates (BFFs) are frequently used in the treatment of progressive systemic pathologies such as osteoporosis or Paget's disease. They are the first choice due to their ability to inhibit osteoclast activity, prevent resorption and modulate bone turnover rate. The plethora of clinical studies that analyses multiple variables to estimate the feasibility of dental implants between users oral BFF, agree that the failure rate is minimal, and generally there are multiple elements, classified into four categories: bacterial factors, factors biomechanical, psychological factors and systemic factors. Its late effects include pain, swelling, paresthesia, discharge, ulcers, fistulas, periodontal pockets, tooth loss, radiolucent osteolytic lesions, among others.

Case report: We documented an exceptional case of the loss of two dental implants osseointegrated and charged rightly, 15 years ago in a 66 years-old patient with osteoporosis who has been taking ibandronic acid orally once a month for the last eight years.

Conclusion: The clinician must cover all preventive and interceptive aspects prior to dental implant treatment in order to prevent failure thereof, despite being low incidence among users of bisphosphonates.

- Oral Presentation 55

TITLE: Hemodynamic monitorization elderly patients during implant procedures

AUTHORS: Gutiérrez Bonet C, García Cubero C, Fernández Cáliz F, Buesa Báñez JM, López-Silva García MJ, Barona Dorado C.

Máster de Cirugía Bucal e Implantología. Hospital Virgen de la Paloma. Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2016 December 15;21(Supplement1):S22.

* doi:10.4317/medoral.17644049

<http://dx.doi.org/10.4317/medoral.17644049>

Aims:

1. To observe the possible alterations of blood pressure (BP), heart rate (HR) and electrocardiogram (ECG) during implant surgery.
2. To study the possible effects on the analyzed constants associated with:
 - Patient's previous pathology
 - A state of anxiety or fear
 - Pain during treatment

Material and Methods: The constants of 100 patients were analyzed, before treatment, at the local anaesthetic

(LA) injection, during treatment, and at the end of it. Anxiety level (Corah test) and experienced pain (VAS) were also evaluated.

Results: Regarding heart rate, 12 tachyarrhythmias, 45 tachycardias and 9 bradyarrhythmias were observed. As for blood pressure, the highest rates were registered at the moment of the injection, 35 patients showed electrocardiographic alterations, 7 in the form of sinus bradycardia, and 28 in the form of tachycardia.

10 patients also showed symptoms, 6 with tachycardia and 4 with dizziness. 9 patients referred discomfort, and regarding anxiety, 26 patients showed moderate anxiety and 12 showed intense anxiety.

Conclusions: Electrocardiographic alterations were present in 35 % of the cases, being sinus tachycardia the most frequent, followed by bradycardia.

1. 48% of the cases showed systolic BP alterations, as well as 42% regarding diastolic pressure. Arterial hypertension was the most frequent alteration and hypotension the least frequent.

2. 10% of the patients showed clinical symptoms

3. Anxiety and hypertension increased the risk of alterations

- Oral Presentation 56

TITLE: Hypophosphatasia. A case report

AUTHORS: Caño Gómez N, Franco Alonso P, Bellarra Arenas C, Franco Carro B, Rubio Alonso L, Martínez González JM.

Máster de Cirugía Bucal e Implantología. Hospital Virgen de la Paloma, Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2016 December 15;21(Supplement1):S22.

* doi:10.4317/medoral.17644050

<http://dx.doi.org/10.4317/medoral.17644050>

Introduction: The hypophosphatasia is a rare inherited disorder characterized by a defect in the mineralization of bone and teeth and a decrease in serum alkaline phosphatase. Depending on when the diagnosis has been made and the severity of the symptoms there are six different clinical forms recognized. It mainly cursed with anormal bone mineralization, premature exfoliation of teeth and severe bone atrophy. The treatment is symptomatic.

The purpose of this communication is to describe the most relevant clinical characteristics of patients with hypophosphatasia with special emphasis on oral repercussions.

Clinical case: Patient, it's a 68 years old woman who comes to our service refereed for rehabilitation of the mandibular arch. She has been diagnosed in adult form of hypophosphatasia without family history and